

## AN ANTIBACTERIAL BIOACTIVE FRACTION OF CINNAMON FRUIT FIELD OF THE INVENTION

The present invention relates to a composition comprising a bioactive fraction obtained from the fruits of *Cinnamomum zeylanicum*, use of the bioactive fraction thus obtained as 5 antibacterial agent and a process for preparing the same from un-conventional parts of *Cinnamomum zeylanicum*.

### PRIOR ART AND BACKGROUND OF THE INVENTION

The problem of food spoilage has plagued man since ancient times. Microbial activity is a primary mode of deterioration of many foods, and the food industry relies heavily upon 10 use of antimicrobial agents to extend shelf life of food products. With growing concern over the presence of chemical residues in foods, the demand for non-toxic natural preservatives has been rising. Natural antibacterial compounds can be used to replace the chemical additives as they have significant antimicrobial activity (Smid EJ, Gorris LGM, Natural antimicrobials for food preservation, In- Shafiurr Rahman M (ed) Handbook of 15 food preservation. Marcel Dekker, Inc, New York, pp.285-308, 1999).

*Cinnamomum zeylanicum* Blume (synonym *Cinnamomum verum* J. S. Presl), the cinnamon of commerce provides various types of oils depending on the part of the plant distilled. Cinnamon is a native of Sri Lanka and tropical Asia. The tree is found in South India up to altitudes of 500 meters but is more common at lower altitudes (*The Wealth of 20 India. A Dictionary of Indian Raw Material and Industrial Products*. Publication and Information Directorate, New Delhi, 1992, p. 582). A total of 53 compounds of cinnamon leaf oil were identified with the major component being eugenol (about 81-84.5%) (Mallavarapu, G.R., Ramesh, S., Chandrasekhara, R.S., Rajeswara Rao, B.R., Kaul, P.N. and Battacharya, A.K. *Flavour and Fragr. J.*, 10, 239, 1995). The essential oil is 25 reported to have antimicrobial (Dubey, N.K. and Mishra, A.K. *Indian Drugs* 27, 529-531, 1990), fungitoxic (Saksena, N.K. and Saksena, S. *Indian Perfumer*, 28, 42-45, 1984), nematicidal (Tiwari, R. Dixit, R. and Dixit, S.N. *Indian Perfumer*, 38, 98-104, 1994), and leech repelling activities (Nakamura, N. Kiuchi, F., Tsuda, Y., Kondo, K. and Sato, T. Shoyakugaka Zasshi, 44, 183-195, 1990). Cinnamaldehyde (about 75%) and camphor 30 (about 56%) have been reported to be the major components of volatile oils from stem bark and root bark respectively, (Senanayake, W.M., Lee, T.H., and wills, R. B. H., *J.Agric. Food. Chem.* 26, 822, 1978). Eugenol and cinnamaldehyde are used as

antimicrobial agents in pharmaceuticals and oral care product (*The Wealth of India. A Dictionary of Indian Raw Material and Industrial Products*. Publication and Information Directorate, New Delhi, 1992, p. 582).

5 Literature survey revealed that, there is no report on the isolation of antibacterial fraction from the fruits of *Cinnamomum zeylanicum*.

The present invention provides a one step process for preparation of a bioactive fraction, which has antibacterial activity. Hence, the inventors have developed a process for the preparation of bioactive fraction from the unconventional parts of *Cinnamomum zeylanicum*, which have no commercial value at present.

10 The principle of the present invention is to provide technology for the preparation of an antibacterial fraction from the unconventional parts of *Cinnamomum zeylanicum*, which can be used as potential natural preservative.

This invention is related to an efficient process for the large-scale preparation of 15 antibacterial fraction from the unconventional parts of *Cinnamomum zeylanicum*.

### **OBJECTS OF THE PRESENT INVENTION**

Therefore, the object of the present invention is to provide a process for the preparation 20 of antibacterial fraction from the fruits of *Cinnamomum zeylanicum*. The process is efficient and it involves the use of simple extraction methods and solvents, which can be re-used.

### **SUMMARY OF THE INVENTION**

According to the present invention, there is provided a composition comprising a 25 bioactive fraction obtained from fruits of *Cinnamomum zeylanicum* having moisture 4-6%, Greenish white colour and mild salty flavor optionally along with one or more pharmaceutically acceptable additives.

The present invention also provides use of the bioactive thus obtained as an antibacterial agent.

The present invention further provides a process for preparing the bioactive fraction.

### **DETAILED DESCRIPTION OF THE INVENTION**

30 Accordingly, the present invention relates to a composition comprising a bioactive fraction obtained from fruits of *Cinnamomum zeylanicum* having

Moisture: 4-6%

Color: Greenish white

Flavor: Mild salty flavor

optionally along with one or more pharmaceutically acceptable additives.

In an embodiment of the present invention the bioactive fraction is a hexane extract  
5 obtained from the fruits of *Cinnamomum zeylanicum*.

In another embodiment of the present invention the composition has antibacterial activity  
against gram positive and gram negative bacterial in the range of 200-500 ppm.

In yet another embodiment of the present invention the composition has antibacterial  
activity against *Bacillus cereus*, *Bacillus subtilis*, *Bacillus coagulans*, *Pseucomonas*  
10 *aeruginosa*, *Staphylococcus aureus*.

The present invention provides a novel use of a bioactive fraction obtained from fruits of  
*Cinnamomum zeylanicum* having

Moisture: 4-6%

Color: Greenish white

15 Flavor: Mild salty flavor

as an antibacterial agent.

In an embodiment of the present invention the bioactive fraction is a hexane extract  
obtained from the fruits of *Cinnamomum zeylanicum*.

In another embodiment of the present invention, the bioactive fraction has antibacterial  
20 activity against gram positive and gram negative bacterial in the range of 200-500 ppm.

In yet another embodiment of the present invention, the bioactive fraction has  
antibacterial activity against *Bacillus cereus*, *Bacillus subtilis*, *Bacillus coagulans*,  
*Pseucomonas aeruginosa*, *Staphylococcus aureus*.

The present invention also provides a process for preparing antibacterial bioactive  
25 fraction having

Moisture: 4-6%

Color: Greenish white

Flavor: Mild salty flavor

from the unconventional parts of *Cinnamomum zeylanicum*, said process comprising the  
30 steps of:

(a) extracting the powdered fruits of *Cinnamomum zeylanicum* with an organic  
solvent at a temperature in the range of 55-60°C for a time period in the range of  
6-8 mesh.

- (b) filtering and concentrating the solvent obtained in step (a) to obtain a concentrate and to recover upto 90% of the solvent;
- (c) drying the concentrate obtained in step (b) in a vacuum oven at 40-50°C under 5 vacuum at 10-25 mm of mercury to obtain the antibacterial bioactive fraction.

In an embodiment of the present invention, the organic solvent used is hexane.

In another embodiment of the present invention, the yield of hexane extract is about 1.5 to 3.0%.

In yet another embodiment of the present invention, the filtration is carried out by 10 conventional methods.

In still another embodiment of the present invention, the concentration temperature is of 55 – 60°C.

In a further embodiment of the present invention, the antibacterial bioactive fraction thus obtained has antibacterial activity against gram positive and gram negative bacterial in 15 the range of 200-500 ppm.

Accordingly, the present invention provides a process for the preparation of antibacterial fraction, which comprises,

- i) Powdering the fruits of *Cinnamomum zeylanicum* to get a particle size 60-80 mesh.
- 20 ii) extracting of the above said material with hexane in a Soxhlet extractor at a temperature of 55-60 °C for a period of 6-8 h.
- iii) filtering the above extract using Whatman filter paper no.1 to obtain the particle free extract.
- iv) distilling the above extract to recover / recycle the solvent up to 90%.
- 25 v) concentrating the above particle free extract at a temperature of 55 - 60 °C
- vi) drying the above concentrated extract using vacuum oven at 40-50 °C under vacuum at 10-25 mm of mercury.
- vii) the product thus obtained had antibacterial activity against different Gram positive and Gram negative bacteria in the range of 200-500 ppm.

30 In an embodiment of the present invention, the yield of hexane extract was found to be 1.5 -3.0%.

The preparation of antibacterial fraction from the unconventional parts of *Cinnamomum zeylanicum* was done according to the flow diagram shown in Figure 1.

The novelty of the process includes:

1. This is the first report of preparation of antibacterial fraction from the unconventional parts of *Cinnamomum zeylanicum*.
- 5 2. The invention is a single step process to obtain the bioactive fraction from the unconventional parts of *Cinnamomum zeylanicum*.

The following examples are given by way of illustration of the present invention and therefore should not be construed to limit the scope of the present invention.

#### Example 1

10 50 g fruits of *Cinnamomum zeylanicum* were powdered using mixer grinder to get a 60 mesh size. The powder was extracted using 200 ml of hexane at 60 °C for 8 h in a Soxhlet extractor. The hexane extract was filtered using Whatman filter paper No.1 and it was concentrated to recover the 150 ml of solvent. The concentrate was dried in a vacuum oven at 40 °C under 10 mm of vacuum. The yield of extract was 1.4 g.

15 The antibacterial assay for the extract of *Cinnamomum zeylanicum* was tested by pour plate method against *Bacillus cereus* by the method of Negi *et al.* (J. Agricultural and Food Chemistry 47, 4297-4300, 1999). To flasks containing 20 ml melted nutrient agar, different concentration of test material in propylene glycol were added. Equivalent amounts of propylene glycol were used as controls. One hundred  $\mu$ l (about  $10^3$  cfu/ml) of

20 culture was inoculated into the flasks under aseptic conditions. The media was then poured into sterilized petri plates in quadruplet and incubated at 37 °C for 20-24 h for growth. The minimum inhibitory concentration (MIC) was reported as the lowest concentration of the compound capable of inhibiting the complete growth of the bacterium being tested. The MIC value of *Cinnamomum zeylanicum* fruit extract against

25 *Bacillus cereus* was 250 ppm.

#### Example -2

The dried fruits (100 g) of *Cinnamomum zeylanicum* were powdered in a mixer grinder to get 80 mesh size. The powder was extracted with 400 ml of hexane by using Soxhlet extractor at 55 °C for 8 h. The extract was filtered using Whatman filter paper No 1. and concentrated under vacuum to recover the 360 ml of solvent. The concentrate dried at a temperature of 35 °C and under a reduced pressure at 25 mm of mercury. The yield of hexane extract was 3.0g.

The antibacterial assay for the extract of *Cinnamomum zeylanicum* was done by known

method. The MIC value of *Cinnamomum zeylanicum* fruit extract against *Bacillus subtilis* was 300 ppm.

#### **Example -3**

5 The dried fruits (150 g) of *Cinnamomum zeylanicum* were powdered in a mixer grinder to get 80 mesh size. The powder was extracted with 600 ml of hexane by using Soxhlet extractor at 55 °C for 8 h. The extract was filtered using Whatman filter paper No 1 and concentrated under vacuum to recover the 520 ml of solvent. The concentrate dried at a temperature of 35 °C and under a reduced pressure at 25 mm of mercury. The yield of  
10 hexane extract was 4.7g.

The antibacterial assay for the extract of *Cinnamomum zeylanicum* was done by known method. The MIC value of *Cinnamomum zeylanicum* extract against *Bacillus coagulans* was 300 ppm.

#### **Example 4**

15 The antibacterial assay for the extract of *Cinnamomum zeylanicum* was done by known method. The MIC value of *Cinnamomum zeylanicum* extract against *Pseudomonas aeruginosa* was 200 ppm.

#### **Example 5**

20 The antibacterial assay for the extract of *Cinnamomum zeylanicum* was done by known method. The MIC value of *Cinnamomum zeylanicum* extract against *Staphylococcus aureus* was 500 ppm.

The advantages of the process are:

1. The process is simple and the solvents used in this process can be regenerated for further use.
- 25 2. The raw material has no commercial value at present.